

Bell correlations in a Bose-Einstein condensate

Roman Schmied¹, Jean-Daniel Bancal^{1,2}, Baptiste Allard¹, Matteo Fadel¹, Valerio Scarani², Philipp Treutlein¹, and Nicolas Sangouard¹

¹*Department of Physics, University of Basel, Switzerland*

²*Centre for Quantum Technologies, National University of Singapore*

The parts of a composite system can share correlations that are stronger than any classical theory allows. These so-called Bell correlations can be confirmed by violating a Bell inequality and represent the most profound departure of quantum from classical physics. We report experiments where we detect Bell correlations between the spins of 480 atoms in a Bose-Einstein condensate [1]. We derive a Bell correlation witness from a recent many-particle Bell inequality [2] involving one- and two-body correlation functions only. Our measurement on a spin-squeezed state [3] exceeds the threshold for Bell correlations by 3.8 standard deviations. Concluding the presence of Bell correlations is unprecedented for an ensemble containing more than a few particles. Our work shows that the strongest possible non-classical correlations are experimentally accessible in many-body systems, and that they can be revealed by collective measurements. This opens new perspectives for using many-body systems in quantum information tasks.

- [1] R. Schmied, J.-D. Bancal, B. Allard, M. Fadel, V. Scarani, P. Treutlein, N. Sangouard, to be published (2016)
- [2] J. Tura, R. Augusiak, A.B. Sainz, T. Vértesi, M. Lewenstein, A. Acín, *Science* 344, 1256 (2014)
- [3] M.F. Riedel, P. Böhi, Y. Li, T.W. Hänsch, A. Sinatra, P. Treutlein, *Nature* 464, 1170 (2010)